

### Alcohol Related Deaths by Sex and Age, Maine, 2001-2005

	Number	Crude Rate		Age-adjusted Rate	
		Rate	95% CI	Rate	95% CI
<b>Total</b>	343	5.3	4.7 - 5.8	<b>4.8</b>	<b>4.3 - 5.4</b>
<b>Gender</b>					
Male	243	7.6	6.7 - 8.7	<b>7.3</b>	<b>6.4 - 8.3</b>
Female	100	3.0	2.4 - 3.6	<b>2.6</b>	<b>2.1 - 3.2</b>
<b>Age (years)</b>					
0 - 19	11	0.7*	0.3 - 1.2	<b>NApp</b>	<b>NApp</b>
20 - 34	44	3.8	2.8 - 5.1	<b>NApp</b>	<b>NApp</b>
35 - 49	84	5.3	4.2 - 6.5	<b>NApp</b>	<b>NApp</b>
50 - 64	88	7.1	5.7 - 8.8	<b>NApp</b>	<b>NApp</b>
65+	111	11.8	9.7 - 14.2	<b>NApp</b>	<b>NApp</b>

Alcohol Related Deaths: an estimate of the total number of deaths attributable to alcohol.

ICD-10 codes: full list of ICD-10 codes available at [http://apps.nccd.cdc.gov/DACH\\_ARDI/Info/ICDCodes.aspx](http://apps.nccd.cdc.gov/DACH_ARDI/Info/ICDCodes.aspx)

Crude rates are deaths per 100,000 population

\*rates based on less than 20 deaths may be unreliable.

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population

95% CI: 95% confidence interval of the rate.

Data Source: Centers for Disease Control and Prevention. Alcohol Related Disease Impact (ARDI) application, 2008. Available at [http://apps.nccd.cdc.gov/DACH\\_ARDI/Default.aspx](http://apps.nccd.cdc.gov/DACH_ARDI/Default.aspx).

NApp = not applicable; age-adjusted rates are not calculated for age-specific groups.